


Human IL-17A protein

Cat #: PRP1052

Size: 5µg/20µg/500µg

Product Information

	Product Name: Human IL-17A protein		
REF	Catalog Number: PRP1052	LOT	Lot Number: Refer to product label
	Purity: > 98 % as determined by SDS-PAGE		
	Storage: Store at -20°C		Preparation method: E. coli
	Shipping: The product is shipped at ambient temperature.		

Background: IL-17A is an effective pro-inflammatory cytokine produced by activated memory T cells. There are at least 6 members of the IL-17 family in humans and mice. Because IL-17A has the same characteristics as IL-1 and TNF- α , it can induce joint inflammation and bone and cartilage destruction. It increases the production of IL-6, induces collagen degradation and reduces collagen synthesis in synovium and cartilage and proteoglycan synthesis in cartilage. IL-17A can also increase bone destruction and reduce its formation. Blocking IL-17A with specific inhibitors provides protective inhibition of cartilage and bone degradation.

Sequence: Amino acid sequence derived from Human IL-17A protein (Ile 20-Ala155) (Q16552) was expressed with 6×His tag at C-terminus.

Protein length: The secreted Human IL-17A protein consists of 136 amino acids and has a predicted molecular mass of 16.5 kDa.

Biological Activity: Measure by its ability to induce IL-6 secretion in 3T3 cells. The ED₅₀ for this effect is <6 ng/mL.


Endotoxin: < 0.01 EU per µg of the protein as determined by the LAL method.

Formulation: Lyophilized from sterile PBS, pH 8.0.

Storage Instructions: Lyophilized protein product should be stored desiccated below -18°C. Upon reconstitution, the protein should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage notes: Always centrifuge tubes before opening. It is recommended to reconstitute the lyophilized recombinant Human

IL-17A protein to a concentration of 0.1-1 mg/mL in sterile ddH₂O, and keep at room temperature for at least 20min to fully dissolve. Please avoid vortex vigorously.



Note: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.