


Human BMP-2 Protein

Cat #: PRP1057

Size: 5µg/20µg/100µg/1mg

Product Information

	Product Name: Human BMP-2 Protein		
REF	Catalog Number: PRP1057	LOT	Lot Number: Refer to product label
	Purity: > 95% as determined by SDS-PAGE		
	Storage: Store at -20°C		Preparation method: E. coli
	Shipping: The product is shipped at ambient temperature.		

Background: BMP-2 protein, like other bone morphogenetic proteins, plays an important role in the development of bone and cartilage. BMP-2 protein is involved in the hedgehog pathway, TGF beta signaling pathway, and cytokine-cytokine receptor interaction. BMP-2 and BMP-7 are osteogenic BMPs that have been demonstrated to potently induce osteoblast differentiation in a variety of cell types. BMP-2, BMP-4 and BMP-7 are known to be of major importance in bone formation and repair. In cancerous tissues BMP-2 protein may play an important role in the progression of glioma.

Sequence: Amino acid sequence derived from Human BMP-2 (Gln283-Arg396)(P12643) was expressed.

Protein length: The recombinant Human BMP-2 consists of 115 amino acids and predicts a molecular mass of 13 kDa.

Biological Activity: Measure by its ability to induce alkaline phosphatase production by ATDC5 cells. The ED₅₀ for this effect is <1.0 µg/mL.


Endotoxin: < 0.1 EU per µg of the protein as determined by the LAL method.

Formulation: The protein was lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate, 0.2 M NaCl, pH 3.5.

Storage Instructions: Lyophilized protein product should be stored desiccated below -18°C. Upon reconstitution, the protein should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (5% HSA, 10% FBS or 0.1% BSA). Please prevent freeze-thaw cycles.

Usage notes: Always centrifuge tubes before opening. It is recommended to reconstitute the lyophilized recombinant Human BMP-2 Protein to a concentration of 0.1-1 mg/mL in sterile ddH₂O, and keep at room temperature for at least 20 min to fully

dissolve. Please avoid vortex vigorously.



Note: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.