


## Human IL-7 Protein, His tag (Animal-Free)

Cat #: PRP1089

Size: 5 µg/20 µg/100 µg/1 mg

### Product Information

	<b>Product Name:</b> Human IL-7 Protein, His tag (Animal-Free)		
<b>REF</b>	<b>Catalog Number:</b> PRP1089	<b>LOT</b>	<b>Lot Number:</b> Refer to product label
	<b>Purity:</b> > 95 % as determined by SDS-PAGE		
	<b>Storage:</b> Store at -20°C		<b>Preparation method:</b> E.coli
	<b>Shipping:</b> The product is shipped at ambient temperature.		

**Background:** Interleukin-7 (IL-7) is a multipotent cytokine belonged to one of the members of IL-2 superfamily. IL-7 has diverse effects on the hematopoietic and immune regulations. IL-7 is a trophic factor that is necessary for both B cell and T cell proliferation and development. In addition, it presents potential antitumor effects in tumors such as glioma, melanoma, lymphoma, leukemia, prostate cancer, and glioblastoma.

**Sequence:** Amino acid sequence derived from Human IL-7 (Asp26-His177) (P13232) was expressed with 6×His tag at the C-terminus.

**Protein length:** The recombinant Human IL-7 Protein consists of 152 amino acids and predicts a molecular mass of 18.3 kDa.

**Biological Activity:** Measured in a cell proliferation assay using PHA-activated human peripheral blood lymphocytes (PBMC). The ED<sub>50</sub> for this effect is <0.8 ng/mL.


**Endotoxin:** <0.1 EU per µg of the protein as determined by the LAL method.

**Formulation:** Lyophilized from sterile PBS, pH 8.0.

**Storage Instructions:** Lyophilized protein product should be stored desiccated below -18°C. Upon reconstitution, the protein should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (5% HSA, 10%FBS or 0.1%BSA). Please prevent freeze-thaw cycles.

**Usage notes:** Always centrifuge tubes before opening. It is recommended to reconstitute the lyophilized recombinant protein to a concentration of 0.1-1 mg/mL using the buffer we provided, and keep at room temperature for at least 20 min to fully

dissolve. Please avoid vortex vigorously.



**Note:** The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.