


Mouse IFN- γ Protein

Cat #: PRP1138

Size: 20 μ g/100 μ g/200 μ g/1mg

Product Information

	Product Name: Mouse IFN- γ Protein		
REF	Catalog Number: PRP1138	LOT	Lot Number: Refer to product label
	Purity: > 98 % as determined by SDS-PAGE		
	Storage: Store at -20°C		Preparation method: E.coli
	Shipping: The product is shipped at ambient temperature.		

Background: IFN- γ is an acid-labile interferon produced by CD4 and CD8 T lymphocytes, as well as activated NK cells. IFN- γ receptors are present in most immune cells, which respond to IFN- γ signaling by increasing the surface expression of class I MHC proteins. This promotes the presentation of antigen to T-helper (CD4+) cells. IFN- γ signaling in antigen-presenting, cells and antigen-recognizing B and T lymphocytes, regulates the antigen-specific phases of the immune response.

Sequence: Amino acid sequence derived from Mouse IFN- γ protein (His23-Cys155)(P01580) with 6 \times His tag at the C-terminus was expressed.

Protein length: The Mouse IFN- γ protein consists of 133 amino acids and has a predicted molecular mass of 16.5 kDa.

Biological Activity: Measure by its ability to anti-viral assay in L-929 cells infected with encephalomyocarditis (EMC) virus. The ED₅₀ for this effect is <0.5 ng/mL. The specific activity of recombinant mouse IFN gamma is approximately >2x 10⁶ IU/mg.


Endotoxin: < 0.01 EU per μ g of the protein as determined by the LAL method.

Formulation: Lyophilized from sterile PBS, pH 7.4.

Storage Instructions: Lyophilized protein product should be stored desiccated below -18°C. Upon reconstitution, the protein should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (5% HSA, 10%FBS or 0.1%BSA). Please prevent freeze-thaw cycles.

Usage notes: Always centrifuge tubes before opening. It is recommended to reconstitute the lyophilized protein to a concentration of 0.1-1 mg/mL in sterile ddH₂O, and keep at room temperature for at least 20 min to fully dissolve. Please

avoid vortex vigorously.



Note: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.