


Mouse IL17A Protein

Cat #: PRP1300

Size: 5 µg/20 µg/100 µg /1 mg

Product Information

	Product Name: Mouse IL17A Protein		
REF	Catalog Number: PRP1300	LOT	Lot Number: Refer to product label
	Purity: > 98 % as determined by SDS-PAGE		
	Storage: Store at -20°C		Preparation method: E.coli
	Shipping: The product is shipped at ambient temperature.		

Background: Interleukin-17 (IL-17) is a potent pro-inflammatory cytokine produced by activated memory T cells. There are at least six members of the IL-17 family in humans and in mice. Mature mouse IL-17A shares 61% and 89% amino acid sequence identity with human and rat IL-17A, respectively. As IL-17 shares properties with IL-1 and TNF-alpha, it may induce joint inflammation and bone and cartilage destruction. This cytokine is found in synovial fluids of patients with rheumatoid arthritis, and produced by rheumatoid arthritis synovium. It increases IL-6 production, induces collagen degradation and decreases collagen synthesis by synovium and cartilage and proteoglycan synthesis in cartilage. IL-17 is also able to increase bone destruction and reduce its formation. Blocking of interleukin-17 with specific inhibitors provides a protective inhibition of cartilage and bone degradation.

Sequence: Amino acid sequence derived from Mouse IL17A (Thr22-Ala158) (Q62386) was expressed.

Protein length: The recombinant Mouse IL17A Protein consists of 137 amino acids and predicts a molecular mass of 15.4 kDa.


Biological Activity: Testing in Progress.

Endotoxin: <0.1 EU per µg of the protein by the LAL method.

Formulation: Lyophilized from 20 mM Tris-HCl, 50 mM NaCl, pH 8.0.

Storage Instructions: Lyophilized protein product should be stored desiccated below -20°C. Upon reconstitution, the protein should be stored at 4°C between 2-7 days and for future use below -20°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage notes: Always centrifuge tubes before opening. It is recommended to reconstitute the lyophilized protein using the buffer we provided not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.



Note: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.